

Procter & Gamble Co.

-PE Product Labels –




RAPID RELIEF
MULTI-SYMPTOM

COLD & FLU
+ CONGESTION

Acetaminophen,
Dextromethorphan HBr,
Phenylephrine HCl

RAPID RELIEF
MULTI-SYMPTOM

COLD & FLU
+ CONGESTION

Acetaminophen,
Diphenhydramine HCl,
Phenylephrine HCl



Headache,
Fever,
Sore Throat,
Minor Aches
& Pains

Nasal Congestion,
Sinus Pressure
Cough

Non-Drowsy

Headache, Fever,
Sore Throat,
Minor Aches
& Pains

Sneezing,
Runny Nose, Cough

Nasal Congestion,
Sinus Pressure

Nighttime Relief

HONEY LEMON
FLAVORED

10 DAYQUIL POWDER PACKETS; 6 NYQUIL POWDER PACKETS; 16 TOTAL POWDER PACKETS

DayQuil (continued)

Directions: • Take only as directed. • Do not exceed 6 doses per 24 hrs. • Dissolve contents of one packet into 8 oz. of hot water and stir briefly or white hot. Consume entire drink within 10-15 minutes.

Warnings: • Do not use if you are taking the blood thinning drug warfarin. • Taking sedatives or tranquilizers may increase drowsiness. • Avoid alcoholic drinks. • Do not use when driving a motor vehicle or operating machinery. • Stop use and ask a doctor if: • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present. • If pregnant or breast-feeding, ask a health professional before use. • Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: • Take only as directed. • Do not exceed 6 doses per 24 hrs. • Dissolve contents of one packet into 8 oz. of hot water and stir briefly or white hot. Consume entire drink within 10-15 minutes.

Warnings: • Do not use if you are taking the blood thinning drug warfarin. • Taking sedatives or tranquilizers may increase drowsiness. • Avoid alcoholic drinks. • Do not use when driving a motor vehicle or operating machinery. • Stop use and ask a doctor if: • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present. • If pregnant or breast-feeding, ask a health professional before use. • Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

NyQuil (continued)

Directions: • Take only as directed. • Do not exceed 6 doses per 24 hrs. • Dissolve contents of one packet into 8 oz. of hot water and stir briefly or white hot. Consume entire drink within 10-15 minutes.

Warnings: • Do not use if you are taking the blood thinning drug warfarin. • Taking sedatives or tranquilizers may increase drowsiness. • Avoid alcoholic drinks. • Do not use when driving a motor vehicle or operating machinery. • Stop use and ask a doctor if: • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present. • If pregnant or breast-feeding, ask a health professional before use. • Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

10 DAYQUIL POWDER PACKETS; 6 NYQUIL POWDER PACKETS; 16 TOTAL POWDER PACKETS

MADE IN MEXICO
DIST. BY: PROCTER & GAMBLE
CINCINNATI, OH 45202

MAX STRENGTH

VICKS

NyQuilTM

HOT REMEDY

RAPID RELIEF MULTI-SYMPTOM

COLD & FLU + CONGESTION

Acetaminophen,
Diphenhydramine HCl,
Phenylephrine HCl



Headache, Fever,
Sore Throat,
Minor Aches
& Pains

Sneezing,
Runny Nose
Cough

Nasal Congestion,
Sinus Pressure

8 POWDER PACKETS

**HONEY
LEMON
FLAVORED**

Nighttime Relief

P&G
www.pg.com
vicks.com
Patents: www.pg.com/patents

Tamper Evident: Do not use
if sachet is opened or torn

Keep outer package
for complete product
information.

MADE IN MEXICO
DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202
90262782

Drug Facts

Active ingredients (in each Packet)	Purpose
Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

• with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- Do not use more than directed
- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information each packet contains:

- potassium 10 mg
- phenylephronics: contains phenylephrine 68 mg per dose
- do not exceed 23°C

Inactive ingredients: acacia, potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions?
1-800-362-1883

U4
239004515
6

PAPER BOX **MULTI-LAYER POUCH**

**Drug Facts****Active ingredients (in each 15 mL)**

	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if
 • adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
 • child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
 • taken with other drugs containing acetaminophen
 • adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema • a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless
 • pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
 • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
 • cough comes back, or occurs with rash or headache that lasts.
 These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed	• only use the dose cup provided	• do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs	
children 6 to under 12 yrs	15 mL every 4 hrs	
children 4 to under 6 yrs	ask a doctor	
children under 4 yrs	do not use	

Other information

• each 15 mL contains: sodium 44 mg
 • Store at no greater than 25°C. Do not refrigerate.

Inactive ingredients

anhydrous citric acid, beta carotene, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions? 1-800-362-1683

TAMPER EVIDENT: Do not use if printed safety seal under cap is missing or damaged.
DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
Made in Canada
 Close cap tightly 90521692



Patents: www.pg.com/patents
www.vicks.com

P&G
www.pg.com

PEEL BACK FOR
 DRUG FACTS



**CLOSE
 TIGHTLY**



Drug Facts (continued)

- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
• adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
• child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
• taken with other drugs containing acetaminophen
• adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin redness • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless

Tamper Evident:
Do not use if printed shrinkband seal around the neck is broken or missing.
← **PEEL BACK FOR DRUG FACTS**

Drug Facts

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - minor aches & pains
 - headache
 - fever
 - sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Dist. by: PROCTER & GAMBLE, CINCINNATI, OH 45202
Patients: www.pg.com/patients
vicks.com

PARENTS:
Learn about how medicine works
www.StopMedicineAbuse.org

91708228

P&G
www.pg.com

PLASTIC BOTTLE

1239000416211 2

**Drug Facts (continued)**

- excitability may occur, especially in children
 - marked drowsiness may occur
 - avoid alcoholic drinks
 - be careful when driving a motor vehicle or operating machinery
 - alcohol, sedatives, and tranquilizers may increase drowsiness
- Stop use and ask a doctor if:**
- you get nervous, dizzy or sleepless
 - pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.
- If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|--|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | do not use unless directed by a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients: citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)**Warnings**

- Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if:
- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
 - child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- thyroid disease
- heart disease
- diabetes
- high blood pressure
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed

Tamper Evident:

Do not use if printed shrinkband seal around the neck is broken or missing.

← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Dorylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headaches
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

DIST. BY: PROCTER & GAMBLE
CINCINNATI, OH 45202
Patents: www.pg.com/patents
vicks.com

PARENTS:
www.StopMedChildUse.org



91765563

P&G

www.pg.com

PLASTIC BOTTLE

3 23900 04161 5



Drug Facts

Active ingredients (in each Packet)

Acetaminophen 300 mg.....Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses Temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores easier breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin redness
- Blisters
- Rash

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease, or for 2 weeks after stopping the MAOI drug). If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed • do not exceed 6 doses per 24 hrs

Age	Dose
adults & children 12 yrs & over	one packet every 4 hours
children under 12 yrs	do not use

• Dissolve contents of one packet into 8 oz. of hot water and stir briskly. Sip while hot. Consume entire drink within 10-15 minutes.
• If using a microwave, add contents of one packet to 8 oz. of cold water, stir briskly before and after heating. Do not overheat.

Other information • each packet contains: potassium 10 mg
• phenylephrine contains phenylalanine 64 mg per dose • do not exceed 25°C

Inactive ingredients: acetylsalicylic acid, aspartame, carboxymethylcellulose, citric acid, DSC Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, leucine, sucrose, tribasic calcium phosphate

Questions? 1-800-362-1683

how2recycle.info

50% MORE

VICKS

50% MORE

VICKS

FluTherapy

SEVERE COLD & FLU

Acetaminophen - Pain reliever/Fever reducer,
Dextromethorphan HBr - Cough suppressant,
Phenylephrine HCl - Nasal decongestant.

• Fast Relief of Cold & Flu symptoms
• Soothing Vicks Vapors

Tamper Evident:
Do not use if inner sealed packet is torn or broken.

MADE IN MEXICO
DIST. BY:
PROCTER & GAMBLE,
CINCINNATI, OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents

6-9
PACKETS

HONEY LEMON FLAVOR

1 2 3 4 5 6 7 8 9
SIDE VIEW OF 9 PACKETS

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Keep outer package for complete product information. 9126-1958

Drug Facts

Active ingredients (in each Packet)	Purpose
Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- Do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- excitability may occur especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs

Age	Dose
Adults & children 12 yrs & over	one packet every 4 hours
Children under 12 yrs	do not use

• Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes.

• If using a microwave, add contents of one packet to 8 oz. of cold water, stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg
- phenylephrine: contains phenylephrine 68 mg per dose
- do not exceed 25°C.

Inactive ingredients: aceulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions? 1-800-362-1683

Keep outer package for complete product information.

91380172



how2recycle.info



50%
MORE



50%
MORE



FluTherapy

SEVERE COLD & FLU

Acetaminophen - Pain reliever/Fever reducer,
Diphenhydramine HCl - Antihistamine/Cough suppressant
Phenylephrine HCl - Nasal decongestant

• Fast Relief of Cold & Flu symptoms
• Soothing Vicks Vapors

Tamper Evident:
Do not use if inner sealed
packet is torn or broken.

MADE IN MEXICO
DIST. BY:
PROCTER & GAMBLE,
CINCINNATI, OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents

69
PACKETS



Delightful & Great
Tasting with Honey
Lemon Flavor.

ACTUAL SIZE



DayQuil

SEVERE+



COLD & FLU

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guafenesin

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Cough
- Chest Congestion

Non-Drowsy

32 DAYQUIL COATED CAPLETS

NyQuil

SEVERE+



COLD & FLU

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

Nighttime Relief

16 NYQUIL COATED CAPLETS, 48 TOTAL COATED CAPLETS

PARENTS:

Keep this and all other medicines out of the reach of children. Use only as directed. Do not use if you are allergic to any of the ingredients. See your doctor if you have a fever that lasts more than 3 days or if you have a cough that lasts more than 10 days.

WARNING: Do not take with other medicines containing acetaminophen. Do not take with other medicines containing phenylephrine. Do not take with other medicines containing dextromethorphan. Do not take with other medicines containing doxylamine. Do not take with other medicines containing guafenesin. Do not take with other medicines containing pseudoephedrine. Do not take with other medicines containing ephedrine. Do not take with other medicines containing phenol. Do not take with other medicines containing salicylates. Do not take with other medicines containing aspirin. Do not take with other medicines containing ibuprofen. Do not take with other medicines containing naproxen. Do not take with other medicines containing acetaminophen. Do not take with other medicines containing phenylephrine. Do not take with other medicines containing dextromethorphan. Do not take with other medicines containing doxylamine. Do not take with other medicines containing guafenesin. Do not take with other medicines containing pseudoephedrine. Do not take with other medicines containing ephedrine. Do not take with other medicines containing phenol. Do not take with other medicines containing salicylates. Do not take with other medicines containing aspirin. Do not take with other medicines containing ibuprofen. Do not take with other medicines containing naproxen.


DayQuil Facts (continued)

Keep this and all other medicines out of the reach of children. Use only as directed. Do not use if you are allergic to any of the ingredients. See your doctor if you have a fever that lasts more than 3 days or if you have a cough that lasts more than 10 days.

NyQuil Facts (continued)

Keep this and all other medicines out of the reach of children. Use only as directed. Do not use if you are allergic to any of the ingredients. See your doctor if you have a fever that lasts more than 3 days or if you have a cough that lasts more than 10 days.

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



DayQuil™ SEVERE+
VICKS VapoCOOL™

COLD & FLU

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guaifenesin

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Cough • Chest Congestion

Non-Drowsy

Drug Facts

Active ingredients (in each caplet)

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion • cough due to minor throat & bronchial irritation
- minor aches & pains • headache • fever • runny nose & sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

2 COATED CAPLETS

USE SCISSORS

Safety sealed: Use only if foil is intact.

Drug Facts (continued)

Allergy Alert: acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use if you have** • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema **Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not use more than directed. Stop use and ask a doctor if** • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed

- do not exceed 8 caplets per 24 hrs
- adults & children 12 yrs & over2 caplets with water every 4 hrs
- children 4 to under 12 yrsask a doctor
- children under 4 yrsdo not use

Other information • each caplet contains: sodium 4 mg

- do not exceed 25°C

Inactive ingredients croscarmellose sodium, crospovidone, FD&C Yellow No. 6 Aluminum Lake, flavor, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions? 1-800-362-1683

P&G
www.pg.com

www.vicks.com

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202 97154736





NyQuilTM SEVERE+TM
VICKS[®] VapoCOOLTM

COLD & FLU

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Nighttime Relief

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose • Cough

Drug Facts
Active ingredients (in each caplet)

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:
 • nasal congestion • cough due to minor throat & bronchial irritation
 • minor aches & pains • headache • fever • runny nose & sneezing

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

2 COATED CAPLETS

FOLD ON DOTTED LINE TO OPEN FOLD & TEAR
 TEAR TOWARDS CENTER



USE SCISSORS

Safety sealed: Use only if foil is intact.

Drug Facts (continued)
Allergy Alert: acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash
 If a skin reaction occurs, stop use and seek medical help right away
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use if you have** • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland **Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers • taking the blood thinning drug warfarin **When using this product • do not use more than directed** • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness **Stop use and ask a doctor if** • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed
 • do not exceed 8 caplets per 24 hrs
 adults & children 12 yrs & over2 caplets with water every 4 hrs
 children 4 to under 12 yrsask a doctor
 children under 4 yrsdo not use

Other information • do not exceed 25°C

Inactive ingredients crospovidone, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions? 1-800-362-1683 www.vicks.com
 DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202 96892693

P&G
www.pg.com

